INTERNATIONAL DEVELOPMENT PLANNING REVIEW
ISSN:1474-6743 | E-ISSN:1478-3401

ANTIBIOTIC USE IN DENTAL CARE DURING PREGNANCY: SAFETY, GUIDELINES, AND EMERGING TRENDS

Ashwag Moohmaed Alahmari1, Sarah Hamed Almutairi2, Alwah Muhammad Alshmrani3, Nouf Ibrahim Almkezem4, Naif Ahmed Alrumayyan5, Ghada Alshahrani6, Munarh Abdullh Alanazi7, Maher Alabdulkarim8, Anwar Yahya Mashi9, Hussam Alshamrani10.

1. Dental assistant, Dental clinic complex west of Riyadh, Saudi Arabia, Ashwag140753@gmail.com
2. Dental hygienist, Dental clinic complex west of Riyadh, Saudi Arabia, Sarah-hamed@hotmail.com
3. Dental assistant, Dental clinic complex west of Riyadh, Saudi Arabia, lwlhalshmrany4@gmail.com
4. Dentist, Druma General Hospital (C3), Saudi Arabia.
5. Dentist, Druma General Hospital (C3), Saudi Arabia.
6. Dental assistant, Dental clinic complex west of Riyadh, Saudi Arabia, Ghabalshahrani@moh.gov.sa
7. Dental assistant, Al Yamamah hospital, Saudi Arabia, Malanazi221@moh.gov.sa
8. Periodontics, Dental clinic complex west of Riyadh, Saudi Arabia, dr.alabdulkarim@gmail.com
9. Dental Assistant, Al Yamamah Hospital, Saudi Arabia, Anmashi@moh.gov.sa
10. Family dentistry specialist, Security forces hospital program Riyadh, Saudi Arabia, dr.hos@live.com

Abstract

Antibiotic use during pregnancy is a critical aspect of dental care, as infections left untreated can lead to severe complications for both the mother and fetus. However, prescribing antibiotics for pregnant patients requires careful consideration to balance the benefits of maternal health against potential risks to fetal development. This article explores the role of antibiotics in managing dental infections during pregnancy, focusing on their indications, safety profiles, and contraindications. Safe antibiotics, such as penicillins, cephalosporins, and clindamycin, are discussed alongside those to avoid, including tetracyclines and quinolones. The article also highlights the risks of teratogenicity, antibiotic resistance, and adverse pregnancy outcomes associated with improper use. Evidence-based guidelines for dentists emphasize patient assessment, trimester-specific considerations, and the importance of collaboration with obstetricians. Emerging trends, such as antibiotic stewardship programs, alternative therapies like photodynamic therapy and probiotics, and advances in personalized medicine, are also explored. By adhering to these principles, dental professionals can manage infections effectively while safeguarding maternal and fetal health.

Introduction

Oral health is a critical component of overall healthcare during pregnancy, as untreated dental infections or diseases can adversely affect both the mother and fetus. Pregnancy induces numerous physiological and hormonal changes that predispose women to oral health issues, including gingivitis, periodontitis, and pulpitis. These conditions, if left untreated, can escalate into serious infections such as odontogenic abscesses or systemic complications, which may necessitate the use of antibiotics (Geevarghese et al., 2017). However, the prescription of antibiotics during pregnancy requires careful consideration to ensure maternal health without compromising fetal safety.

The management of dental infections during pregnancy is particularly challenging due to concerns about teratogenic risks, alterations in drug pharmacokinetics, and the potential for adverse pregnancy outcomes. The American Dental Association (ADA) and the American College of Obstetricians and Gynecologists (ACOG) emphasize the importance of providing timely and appropriate dental care during pregnancy, including the use of antibiotics when clinically indicated (Aliabadi et al., 2022). Antibiotics play a vital role in managing odontogenic infections, preventing systemic spread, and minimizing complications. However, their use must be guided by evidence-based practices to balance therapeutic benefits with potential risks.

Certain antibiotics, such as penicillins, cephalosporins, and clindamycin, have been deemed safe for use during pregnancy, while others, such as tetracyclines and quinolones, are contraindicated due to their association with fetal toxicity and developmental issues (Cohlan, 1977; Ziv et al., 2018). Despite the availability of guidelines, studies reveal variability in prescribing practices among dental professionals, often influenced by a lack of knowledge or awareness about safe antibiotic use in pregnancy (Ibhawoh & Enabulele, 2015).

This article explores the role of antibiotics in dental care for pregnant women, highlighting their indications, safety profiles, and potential risks. By providing evidence-based insights, it aims to equip dental professionals with the knowledge to make informed decisions, ensuring optimal care for pregnant patients while safeguarding fetal health.

1. Oral Health Challenges During Pregnancy

Pregnancy brings about significant hormonal, physiological, and behavioral changes that can affect a woman's oral health. These changes often exacerbate pre-existing conditions or lead to the development of new oral health problems, making pregnant women more susceptible to dental complications. Common oral health issues during pregnancy include gingivitis, periodontitis, dental caries, and pulpitis, which can negatively impact both maternal and fetal health if left untreated (Geevarghese et al., 2017).

1.1 Hormonal Changes and Gingival Health

The hormonal fluctuations during pregnancy, particularly elevated levels of estrogen and progesterone, increase blood flow to the gums and heighten the inflammatory response to plaque. This leads to pregnancy gingivitis, characterized by swollen, red, and bleeding gums, affecting up

to 75% of pregnant women (Sant'Ana et al., 2011). If untreated, gingivitis can progress to periodontitis, a more severe condition involving the destruction of the supporting structures of teeth.

1.2 Periodontal Disease and Adverse Pregnancy Outcomes

Periodontitis has been associated with adverse pregnancy outcomes such as preterm birth, low birth weight, and preeclampsia. The systemic inflammatory response triggered by periodontal infections may lead to increased levels of inflammatory mediators, which are implicated in preterm labor (Sant'Ana et al., 2011; Romero et al., 2001). Therefore, addressing periodontal health during pregnancy is crucial not only for oral health but also for overall pregnancy outcomes.

1.3 Increased Risk of Dental Caries

Behavioral changes, such as dietary cravings for sugary foods and decreased attention to oral hygiene due to nausea or fatigue, contribute to an increased risk of dental caries during pregnancy (Hashim, 2012). Untreated caries can lead to pulpitis and odontogenic infections, which may require antibiotic therapy or even surgical intervention.

1.4 Pulpitis and Dental Pain

The increased vascularization and hormonal changes during pregnancy may also exacerbate pulp sensitivity, leading to pulpitis. Acute odontogenic infections resulting from untreated pulpitis can progress to abscess formation, cellulitis, or systemic complications, which can pose risks to both maternal and fetal health (Aliabadi et al., 2022).

1.5 Barriers to Dental Care

Despite the importance of oral health, many pregnant women fail to seek timely dental care due to misconceptions about the safety of dental procedures during pregnancy or a lack of access to care (Marchi et al., 2010). Studies show that a significant proportion of pregnant women experience untreated dental issues, which can escalate into more severe complications if neglected.

Addressing oral health challenges during pregnancy is critical to ensuring both maternal and fetal well-being. Early diagnosis and treatment, including the appropriate use of antibiotics in cases of infection, are essential for managing oral health problems effectively and preventing complications.

2. The Role of Antibiotics in Dental Care for Pregnant Women

Antibiotics are used in dentistry to treat or prevent bacterial infections associated with endodontic and periodontal diseases. In pregnant patients, however, antibiotic therapy must be approached cautiously due to the potential for drug teratogenicity and adverse fetal effects (Aliabadi et al., 2022). Antibiotics are indicated in the following situations:

 Odontogenic Infections: Severe infections, such as abscesses, cellulitis, and periapical infections, require antibiotics to control the spread of bacteria and prevent systemic complications (Ibhawoh & Enabulele, 2015).
 Prophylaxis for High-Risk Patients: Antibiotics are prescribed prophylactically for patients at risk of infective endocarditis or those undergoing invasive dental procedures, including extractions and periodontal surgeries (Little et al., 2007).

 Acute Periodontal Infections: Periodontal infections in pregnant women are associated with systemic inflammation, which can increase the risk of adverse pregnancy outcomes, such as preterm birth and low birth weight. Antibiotics may be necessary to control the infection and reduce systemic inflammation (Sant'Ana et al., 2011).

3. Safe Antibiotics During Pregnancy

The prescription of antibiotics during pregnancy must strike a balance between effectively managing maternal infections and ensuring fetal safety. While some antibiotics are considered safe for use during pregnancy, others pose risks of teratogenicity, developmental abnormalities, or adverse pregnancy outcomes. The safety of antibiotics is often determined by their pharmacokinetic properties, ability to cross the placental barrier, and their effects on fetal development. This section discusses antibiotics deemed safe for pregnant women based on current research and clinical guidelines.

3.1 Penicillins

Penicillins, including amoxicillin and ampicillin, are widely considered the safest antibiotics during pregnancy. They are effective against a broad spectrum of gram-positive and gram-negative bacteria and have minimal teratogenic risk.

- **Safety Profile**: Penicillins are classified as Pregnancy Category B by the FDA, indicating no evidence of harm to the fetus in animal studies or limited human data (Lin et al., 2012).
- **Clinical Indications**: Amoxicillin is frequently prescribed for odontogenic infections, such as periapical abscesses and periodontitis, due to its well-established efficacy and safety record.

3.2 Cephalosporins

Cephalosporins, such as ceftriaxone and cefazolin, are considered safe alternatives for pregnant women who are allergic to penicillins. They are structurally similar to penicillins but have a broader antibacterial spectrum.

- **Safety Profile**: Like penicillins, cephalosporins are classified as Pregnancy Category B. They show no significant association with congenital anomalies (Czeizel et al., 2001).
- **Clinical Indications**: Cephalosporins are commonly used for infections requiring broader coverage, such as cellulitis, odontogenic infections, and post-surgical prophylaxis.

3.3 Clindamycin

Clindamycin is an effective antibiotic for managing anaerobic bacterial infections, including periodontal and endodontic infections. It is particularly useful for patients with penicillin allergies.

- **Safety Profile**: Clindamycin is classified as Pregnancy Category B and has been extensively studied, showing no evidence of fetal harm (Lamont et al., 2011).
- **Clinical Indications**: Clindamycin is often prescribed for periodontal abscesses, osteomyelitis, and severe odontogenic infections.

3.4 Metronidazole

Metronidazole is an effective antibiotic for anaerobic infections and is often used in combination with other antibiotics to treat severe dental infections.

- **Safety Profile**: Although some controversy surrounds metronidazole, studies suggest that its use in the second and third trimesters does not increase the risk of congenital anomalies (Burtin et al., 1995). However, its use during the first trimester should be avoided unless absolutely necessary.
- **Clinical Indications**: Metronidazole is effective for treating acute necrotizing ulcerative gingivitis (ANUG), periodontitis, and other anaerobic infections.

3.5 Erythromycin

Erythromycin, a macrolide antibiotic, is another safe option during pregnancy, particularly for patients allergic to penicillins.

- **Safety Profile**: Erythromycin is classified as Pregnancy Category B. However, care must be taken to avoid the estolate form, as it has been associated with hepatotoxicity during pregnancy (McCormack et al., 1977).
- **Clinical Indications**: Erythromycin is used for infections caused by gram-positive bacteria, as well as periodontal abscesses.

3.6 Azithromycin

Azithromycin, another macrolide, is preferred for its shorter treatment duration and broader spectrum of activity.

- **Safety Profile**: Azithromycin is classified as Pregnancy Category B and is considered safe based on current research (Romøren et al., 2012).
- **Clinical Indications**: It is effective against soft tissue infections, periodontal infections, and odontogenic abscesses.

3.7 Considerations for Safe Use

While the above antibiotics are generally safe, their use should be guided by clinical judgment and evidence-based guidelines:

1. **Trimester Considerations**: The timing of antibiotic use is crucial. Most antibiotics are safest during the second and third trimesters, as the risks of teratogenicity are higher during the first trimester when organogenesis occurs.
2. **Dosage and Duration**: Prescribing the correct dosage and duration is essential to minimize risks. Overuse or misuse of antibiotics can contribute to antibiotic resistance and adverse maternal or fetal outcomes (Aliabadi et al., 2022).
3. **Medical History**: A thorough review of the patient's medical history, including allergies and previous antibiotic use, is essential to avoid adverse reactions.

4. Antibiotics to Avoid During Pregnancy

While antibiotics play a critical role in managing dental infections during pregnancy, certain classes of antibiotics are known to pose significant risks to fetal development and are generally contraindicated. The safety of these drugs is often compromised due to their teratogenic potential, adverse effects on fetal growth, or long-term developmental consequences. This section discusses antibiotics that should be avoided during pregnancy, their associated risks, and the rationale for their exclusion in the management of pregnant patients.

4.1 Tetracyclines

Tetracyclines, including doxycycline and minocycline, are among the most widely recognized contraindicated antibiotics during pregnancy.

- **Risks to Fetal Development**: Tetracyclines are known to cross the placental barrier and accumulate in developing fetal bones and teeth. This can result in permanent discoloration of primary and secondary teeth, hypoplasia of dental enamel, and inhibition of fetal bone growth (Cohlan, 1977; Cohlan et al., 1963).
- **Teratogenic Effects**: Studies have shown that tetracyclines can interfere with skeletal development, particularly during the second and third trimesters when fetal mineralization is at its peak.
- **Clinical Recommendation**: Tetracyclines should be strictly avoided throughout pregnancy, as safer alternatives like penicillins or cephalosporins are available (Aliabadi et al., 2022).

4.2 Quinolones

Quinolones, such as ciprofloxacin and levofloxacin, are broad-spectrum antibiotics that are generally avoided during pregnancy due to potential adverse effects on fetal cartilage development.

- **Risks to Fetal Development**: Animal studies have linked quinolones to cartilage damage and arthropathy, raising concerns about their impact on fetal joint and skeletal development (Ziv et al., 2018). Human studies, though limited, suggest a similar risk.
- **Clinical Recommendation**: Quinolones should not be prescribed during pregnancy unless no safer alternatives are available, as their potential to harm fetal musculoskeletal development outweighs the benefits.

4.3 Sulfonamides

Sulfonamides, such as sulfamethoxazole, are another class of antibiotics to be avoided, particularly in the first trimester and near term.

- **Risks of Teratogenicity**: Sulfonamides are associated with a higher risk of congenital anomalies, including neural tube defects and cardiovascular abnormalities, when used during the first trimester (Muanda et al., 2017).
- **Neonatal Risks**: Near term, sulfonamides can cause kernicterus (bilirubin-induced brain damage) in the neonate due to their ability to displace bilirubin from albumin-binding sites.
- **Clinical Recommendation**: Safer alternatives such as cephalosporins or macrolides should be considered instead of sulfonamides.

4.4 Aminoglycosides

Aminoglycosides, including gentamicin and streptomycin, are avoided during pregnancy due to their potential for ototoxicity and nephrotoxicity.

- **Risks to Fetal Development**: Aminoglycosides can cross the placenta and accumulate in fetal tissues, particularly the kidneys and inner ear, leading to permanent hearing loss (ototoxicity) and kidney damage (nephrotoxicity) (Aliabadi et al., 2022).
- **Clinical Recommendation**: Aminoglycosides should only be used in life-threatening infections when no other safer options are available, and their use must be closely monitored.

4.5 Chloramphenicol

Chloramphenicol is another antibiotic contraindicated during pregnancy due to its severe adverse effects on the fetus.

- **Risks to Fetal Development**: Chloramphenicol can cause "gray baby syndrome" in neonates, a potentially fatal condition characterized by cyanosis, cardiovascular collapse, and respiratory distress. This occurs because fetal liver enzymes are immature and unable to metabolize the drug effectively (Burtin et al., 1995).
- **Clinical Recommendation**: Chloramphenicol should not be used during pregnancy, as its risks far outweigh its benefits.

4.6 Nitrofurantoin (Near Term)

Although nitrofurantoin is generally considered safe for use during most of pregnancy, it is contraindicated near term (after 36 weeks) due to its potential to cause hemolytic anemia in neonates with glucose-6-phosphate dehydrogenase (G6PD) deficiency (Aliabadi et al., 2022).

4.7 Other Antibiotics with Limited Data

Certain antibiotics, such as rifampin and vancomycin, lack sufficient safety data for use during pregnancy. While they may be used in specific, life-threatening infections, their risks and benefits must be weighed carefully.

5. Risks and Considerations

The use of antibiotics during pregnancy, while often necessary to manage dental infections, comes with potential risks that require careful consideration. These risks extend beyond the immediate therapeutic effects on the mother to include potential adverse outcomes for the developing fetus. Dentists must weigh the benefits of antibiotic therapy against the risks to ensure the safety of both the mother and the baby. This section highlights the key risks and considerations involved in prescribing antibiotics during pregnancy.

5.1 Teratogenic Effects

The primary concern with antibiotic use during pregnancy is the risk of teratogenicity, which refers to the potential of certain medications to cause congenital abnormalities in the fetus. Some antibiotics, such as tetracyclines and quinolones, have been shown to interfere with fetal development. Tetracyclines, for example, can cause permanent discoloration of teeth and impaired bone growth when used during the second or third trimesters (Cohlan, 1977), while quinolones are associated with cartilage and musculoskeletal damage in animal studies (Ziv et al., 2018).

5.2 Adverse Pregnancy Outcomes

Inappropriate or excessive use of antibiotics during pregnancy has been linked to adverse outcomes, including spontaneous abortion, preterm labor, and low birth weight. For example, studies have found that sulfonamides and nitrofurantoin, when used near term, may increase the risk of neonatal complications such as kernicterus and hemolytic anemia (Muanda et al., 2017). These risks underline the importance of prescribing antibiotics only when clinically necessary and avoiding those with known risks.

5.3 Antibiotic Resistance

The misuse or overprescription of antibiotics contributes to the growing global issue of antimicrobial resistance. Pregnant women treated with unnecessary antibiotics for minor or non-bacterial infections can develop resistant bacterial strains, which may complicate future treatments (Koyuncuoglu et al., 2017). This highlights the need for evidence-based prescribing practices to ensure antibiotics are used judiciously.

5.4 Drug Interactions

Pregnancy alters drug metabolism, which can affect the efficacy and safety of antibiotics. Antibiotics metabolized by the liver, such as erythromycin, may pose a higher risk of hepatotoxicity in pregnant women (McCormack et al., 1977). Similarly, antibiotics like metronidazole or clindamycin may interact with other medications the patient is taking, such as prenatal vitamins or anticoagulants, necessitating careful review of the patient's medical history.

5.5 Fetal and Neonatal Health

The timing of antibiotic use during pregnancy is critical. Drugs used during the first trimester, when organogenesis occurs, carry the highest risk of teratogenicity. Meanwhile, antibiotics administered in the later stages of pregnancy, such as nitrofurantoin, can affect neonatal health, particularly in cases of glucose-6-phosphate dehydrogenase (G6PD) deficiency (Aliabadi et al., 2022).

6. Guidelines for Dentists

The prescription of antibiotics for pregnant patients requires adherence to strict guidelines to ensure both maternal health and fetal safety. Dentists play a crucial role in managing oral health during pregnancy and must carefully evaluate the clinical scenario, the patient's medical history, and the risks associated with antibiotic use. The following guidelines provide evidence-based recommendations to help dental professionals make informed decisions when treating pregnant patients.

6.1 Comprehensive Patient Assessment

Before prescribing antibiotics, a thorough assessment of the patient's medical and obstetric history is essential. Dentists should collaborate with the patient's obstetrician to determine any potential risks or contraindications. Factors such as the stage of pregnancy, comorbid conditions (e.g., diabetes, hypertension), and current medications must be considered to avoid adverse effects or drug interactions (Giglio et al., 2009).

6.2 Indications for Antibiotic Use

Antibiotics should only be prescribed when there is a clear clinical indication. Conditions warranting antibiotic use include severe odontogenic infections, abscesses, cellulitis, and systemic signs of infection such as fever or lymphadenopathy. Prophylactic antibiotics may also be necessary for high-risk patients, such as those with a history of infective endocarditis or undergoing invasive dental procedures (Little et al., 2007).

6.3 Choosing the Right Antibiotic

Dentists should prioritize antibiotics with well-established safety profiles during pregnancy. Penicillins, cephalosporins, clindamycin, and erythromycin (without the estolate form) are considered safe options for managing dental infections in pregnant patients (Aliabadi et al., 2022).

Conversely, antibiotics like tetracyclines, quinolones, and sulfonamides should be avoided due to their potential teratogenic effects and risks to fetal development (Cohlan, 1977; Ziv et al., 2018).

6.4 Trimester-Specific Considerations

The timing of antibiotic use is critical. During the first trimester, when organogenesis occurs, caution must be exercised to avoid potential teratogenic effects. Antibiotics deemed safer are optimal choices during this period. The second trimester is generally the safest time for dental interventions requiring antibiotics, while the third trimester requires attention to drugs that may affect neonatal health, such as those causing kernicterus or hemolysis (Muanda et al., 2017).

6.5 Adherence to Dosage and Duration

Dentists must prescribe the appropriate dosage and duration of antibiotics to ensure efficacy while minimizing risks. Overprescription or inadequate treatment can lead to antibiotic resistance or incomplete resolution of the infection (Koyuncuoglu et al., 2017). Patients should be advised to complete the full course of antibiotics even if symptoms subside.

6.6 Patient Education and Communication

Educating pregnant patients about the importance of oral health and the safe use of antibiotics is essential. Clear instructions should be provided regarding antibiotic use, potential side effects, and the importance of adhering to prescribed dosages. Dentists should also address any concerns or misconceptions patients may have about the safety of dental procedures and medications during pregnancy (Silk et al., 2008).

7. Emerging Trends and Future Directions

The management of dental infections in pregnant women is evolving with the advancement of research and technology, emphasizing safer and more effective approaches for both the mother and fetus. Emerging trends and future directions focus on improving antibiotic stewardship, exploring alternative therapies, and integrating personalized medicine into maternal oral healthcare.

7.1 Antibiotic Stewardship

Antibiotic resistance remains a global concern, prompting efforts to minimize unnecessary antibiotic use in dental practice, particularly for pregnant patients. Antibiotic stewardship programs aim to promote evidence-based prescribing practices, ensuring antibiotics are used only when clinically necessary and in appropriate doses and durations. These programs also encourage the use of narrow-spectrum antibiotics when possible, reducing the risk of resistance (Machowska et al., 2020).

7.2 Alternative and Adjunctive Therapies

Non-antibiotic therapies are gaining attention as safer alternatives or adjuncts to traditional treatments for dental infections in pregnant women. For example:

- **Photodynamic Therapy (PDT):** This approach uses light-activated compounds to destroy bacteria in periodontal infections without the need for antibiotics (Ather et al., 2020).
- **Probiotics:** Emerging evidence suggests that probiotics can help control oral microbial imbalances, reducing the risk of infections without harming the fetus.

7.3 Personalized Medicine

Advances in genetic research and pharmacogenomics could pave the way for personalized medicine in dental care. Identifying genetic predispositions to conditions like antibiotic hypersensitivity or resistance can help tailor treatments to individual patients, improving safety and efficacy.

7.4 Increased Research Focus

Future research should focus on long-term studies of antibiotic safety during pregnancy, particularly for less-studied drugs. Additionally, improving public and professional awareness about the importance of maternal oral health and safe antibiotic use is crucial for enhancing outcomes (Aliabadi et al., 2022).

Conclusion

Managing dental infections during pregnancy involves a delicate balance between safeguarding the health of the mother and minimizing risks to the developing fetus. Antibiotics play a critical role in treating infections, but their use must be guided by evidence-based principles. Safe options, such as penicillins, cephalosporins, and clindamycin, should be prioritized, while contraindicated drugs like tetracyclines and quinolones must be avoided. Dentists should carefully assess the patient's medical history, consider trimester-specific risks, and collaborate with obstetricians to ensure appropriate care. The misuse of antibiotics can lead to complications such as teratogenic effects, adverse pregnancy outcomes, and antibiotic resistance, highlighting the importance of rational prescribing practices. Emerging trends, such as alternative therapies and personalized medicine, offer promising avenues for safer and more effective management of dental infections in pregnant women. By adhering to these guidelines, dental professionals can optimize care for pregnant patients, ensuring both maternal and fetal well-being.

References

Aliabadi, T., Saberi, E. A., Motameni Tabatabaei, A., & Tahmasebi, E. (2022). Antibiotic use in endodontic treatment during pregnancy: A narrative review. *European Journal of Translational Myology, 32*(4), 10813. https://doi.org/10.4081/ejtm.2022.10813

Ather, A., Zhong, S., Rosenbaum, A. J., Quinonez, R. B., & Khan, A. A. (2020). Pharmacotherapy during pregnancy: An endodontic perspective. *Journal of Endodontics, 46*(9), 1185–1194. https://doi.org/10.1016/j.joen.2020.05.024

Burtin, P., Taddio, A., Ariburnu, O., Einarson, T. R., & Koren, G. (1995). Safety of metronidazole in pregnancy: A meta-analysis. *American Journal of Obstetrics and Gynecology, 172*(2 Pt 1), 525–529. https://doi.org/10.1016/0002-9378(95)90567-7

Cengiz, S. B. (2007). The pregnant patient: Considerations for dental management and drug use. *Quintessence International, 38*(3), e133–e142.

Cohlan, S. Q. (1977). Tetracycline staining of teeth. *Teratology, 15*(1), 127–129. https://doi.org/10.1002/tera.1420150117

Czeizel, A. E., Rockenbauer, M., Sørensen, H. T., & Olsen, J. (2001). Use of cephalosporins during pregnancy and in the presence of congenital abnormalities: A population-based, case-control study. *American Journal of Obstetrics and Gynecology, 184*(6), 1289–1296. https://doi.org/10.1067/mob.2001.113905

Geevarghese, A., Baskaradoss, J. K., & Sarma, P. S. (2017). Oral health-related quality of life and periodontal status of pregnant women. *Maternal and Child Health Journal, 21*(8), 1634–1642. https://doi.org/10.1007/s10995-016-2255-y

Giglio, J. A., Lanni, S. M., Laskin, D. M., & Giglio, N. W. (2009). Oral health care for the pregnant patient. *Journal of the Canadian Dental Association, 75*(1), 43–48.

Ibhawoh, L., & Enabulele, J. (2015). Endodontic treatment of the pregnant patient: Knowledge, attitude and practices of dental residents. *Nigerian Medical Journal, 56*(5), 311–316. https://doi.org/10.4103/0300-1652.170386

Koyuncuoglu, C. Z., Aydin, M., Kirmizi, N. I., Aydin, V., Aksoy, M., Isli, F., & Akici, A. (2017). Rational use of medicine in dentistry: Do dentists prescribe antibiotics in appropriate indications? *European Journal of Clinical Pharmacology, 73*(8), 1027–1032. https://doi.org/10.1007/s00228-017-2258-7

Lamont, R. F., Nhan-Chang, C. L., Sobel, J. D., Workowski, K., Conde-Agudelo, A., & Romero, R. (2011). Treatment of abnormal vaginal flora in early pregnancy with clindamycin for the prevention of spontaneous preterm birth: A systematic review and meta-analysis. *American Journal of Obstetrics and Gynecology, 205*(3), 177–190. https://doi.org/10.1016/j.ajog.2011.03.047

Lin, K. J., Mitchell, A. A., Yau, W. P., Louik, C., & Hernández-Díaz, S. (2012). Maternal exposure to amoxicillin and the risk of oral clefts. *Epidemiology, 23*(5), 699–705. https://doi.org/10.1097/EDE.0b013e318258cb05

Little, J. W., Falace, D., Miller, C., & Rhodus, N. L. (2007). *Dental Management of the Medically Compromised Patient*. Elsevier Health Sciences.

Machowska, A., Sihavong, A., Eriksen, J., Vongsouvath, M., Marrone, G., & Lundborg, C. S. (2020). Containment of antibiotic resistance measures to improve antibiotic use in pregnancy, childbirth, and young children (CAREChild): A protocol. *BMJ Open, 10*(11), e040334. https://doi.org/10.1136/bmjopen-2020-040334

Muanda, F. T., Sheehy, O., & Bérard, A. (2017). Use of antibiotics during pregnancy and the risk of major congenital malformations: A population-based cohort study. *British Journal of Clinical Pharmacology, 83*(11), 2557–2571. https://doi.org/10.1111/bcp.13364

Sant'Ana, A. C., Campos, M. R., Passanezi, S. C., Rezende, M. L., Greghi, S. L., & Passanezi, E. (2011). Periodontal treatment during pregnancy decreases the rate of adverse pregnancy outcomes: A controlled clinical trial. *Journal of Applied Oral Science, 19*(2), 130–136. https://doi.org/10.1590/s1678-77572011000200009

Ziv, A., Masarwa, R., Perlman, A., Ziv, D., & Matok, I. (2018). Pregnancy outcomes following exposure to quinolone antibiotics: A systematic review and meta-analysis. *Pharmaceutics Research, 35*(5), 109. https://doi.org/10.1007/s11095-018-2383-8